

From the Labs to the Clinics

Published April 8th, 2026

Robert V. Harrison, PhD, DSc

In Praise of Noise Dosimetry

We have seen much “new knowledge” that can potentially impact hearing healthcare. For example, haircell growth from stem-cells, auditory nerve regeneration, gene therapy, the role of AI in hearing prosthesis development and employment, etc. I have covered some of these areas in previous “Labs to Clinics” columns. Most of these new developments relate to possible hearing restoration. Today, however, rather than discuss possible new ways to treat hearing loss, I want to revisit some existing concepts and methods in audiology for the **prevention** of noise-induced hearing loss. For many years I was involved in the prevention of noise-induced hearing loss in children, particularly education campaigns about the safe use of personal sound entertainment devices, originally cassette and MP3 players, more recently smartphones and videogame devices. See Figure 1 below [1, 2]. For the most part, we are (I hope) much more aware about the risks of exposure to loud acoustic signals.

www.soundsense.ca

Save your hearing for the music



IT'S TOO LOUD when:
you have to shout to be heard, or when you're wearing headphones, you can't hear your friends talking to you.

Use some Sound Sense.
Enjoy yourself but save your hearing for the music. Turn down the volume, wear ear plugs at loud events (rolled up tissue doesn't help!) and keep away from the speakers.

Look out for the Warning Signs of hearing damage!

- Your ears "ring" after a dance, concert or playing music
- You're turning the TV up louder to understand it
- You can't understand your friends if there's background noise
- You need to sit close to the teacher to hear
- Everyone seems to mumble or talk too quickly

(If you have any of the above symptoms, tell a parent or your teacher.)

Decibel Level	How long you can listen with unprotected hearing	Noise Source
130+	Beyond pain threshold	Boom car, jet takeoff, jackhammer
120	Danger!	Dance club
115	Under 1 minute	Rock concert, sports arena
109	Under 2 minutes	Car horn, leaf blower, power saw
106	3.75 minutes	Portable CD/mp3 player (at high volume)
103	7.5 minutes	Belt sander
100	15 minutes	School dance, snowmobile, factory machines
97	30 minutes	Wet lay, motorcycle
94	1 hour	Electric drill
91	2 hours	Shedding, blender, tractor, lawn mower
▲ LEVEL AT WHICH DAMAGE BEGINS ▲		
85	8 hours	Vacuum cleaner, hand saw
75	Safe	Conversation

hearing foundation of canada

Figure 1. Sound Sense poster from The Hearing Foundation of Canada, targeting young people with the slogan “Save your hearing for the music”. [1,2]

In the broader field of environmental and occupational noise, I have come to appreciate the role of Occupational Health and Safety (OHS) specialists (often together with Audiologists) and their methods of noise dosimetry. The basic concepts of noise exposure dosimetry are not new, having been established soon after the Second World War. This methodology has been most instrumental in limiting harmful noise exposure and establishing safe working conditions (often backed by legislative guidelines). Noise dosimetry deserves some good press!

We all understand that acoustic trauma is not the same as noise-induced hearing loss (NIHL), although I have often seen the terms used interchangeably. Perhaps there are some formal definitions, but for me it is simple. If an acoustic signal is at a pressure level that can physically injure the ear(s), and cause an immediate and permanent hearing problem, then it is acoustic trauma. At the less severe end of the scale, a much lower-intensity noise exposure over long periods can lead to hearing problems. This is how I would characterize noise-induced hearing loss. Here, the hearing loss risk depends on both signal intensity and duration. A relatively intense, 30-second acoustic signal can have the same risk as 8 hours of continuous low-level noise.

If you are in an environment where the noise level feels excessive, it is routine to consult the sound level app on your phone (no self-respecting audiologist is without the app). The revealed level may be uncomfortable (as in “let’s get out of this restaurant”), but it will not definitively indicate risk of NIHL without **an exposure duration**. The intensity/time trade-off is the key feature about noise

dosimetry.

Noise exposure dosimetry involves continuous monitoring of acoustic signal levels (usually A-weighted) using a microphone attached to the body, typically at shoulder level, together with a data storage device. In the occupational health and safety world, this monitoring is often conducted over a full working shift or a long-term period in a representative sound environment. Most often the goal is to measure a long-term average exposure level over an 8-hour workday (in any 24-hour period). Based on hundreds of published studies, the daily noise dose that can pose a risk of hearing loss has been established and formally adopted by health and safety regulatory bodies. For example, we have the Canadian Centre for Occupational Health and Safety (CCOHS) [4] the Occupational Safety and Health Administration (OSHA) in the US [3]. There are multiple jurisdictions worldwide, but all have set similar guidelines about the length of time that a person can be exposed to various levels of continuous noise before there is a risk to hearing. This is set at 85-87 dBA for an 8-hour time-weighted average (TWA) over any 24-hour period. This level is referred to as an action level at which some mitigation procedure must be implemented, often requiring the mandatory use of hearing protectors. In Canada this risk threshold (action level) is set at 87 dBA TWA.

Table 1 below is based on Canadian Centre for Occupational Health and Safety guidelines [ref 4] and shows the relationship between noise exposure level and permissible exposure time. I have indicated on this table the relevant risk threshold of 87 dBA for an 8hr daily exposure.

The below table demonstrates the relationship between sound level exposure and permissible exposure in hours. As per the table, the regulations allow for up to sixteen (16) hours of 84 dBA in any 24 hour period, and the maximum permitted exposure level for eight (8) hours is 87 dBA.

Maximum duration of exposure to a-weighted sound pressure levels in the work place

A-weighted sound pressure level (dBA)	Maximum duration of exposure in hours/minutes per employee per 24-hour period
84	16.0
87	8.0
90	4.0
93	2.0
96	1.0
99	30 minutes
102	15 minutes
105	7.5 minutes
108	3.75 minutes
111	<2 minutes
114	<1 minute

TABLE 1. Based Canadian Centre for Occupational Health and Safety (2025). Noise, occupational Exposure limits in Canada [4].

To reiterate, the important concept in dosimetry is the trade-off between noise exposure level and duration. Table 1 shows a range of long-term average noise levels (left column) versus exposure duration (right) that can pose a risk of noise-induced hearing loss. The table is a useful reference guide to many more noise exposure situations than just the occupational “working day”. The table

indicates the risk thresholds for noise exposure across various signal intensities and durations.

Indeed, there are numerous useful applications of noise dosimetry beyond the formal workplace. There are dosimetry studies of road traffic noise, train and airport noise, construction-site noise, entertainment venue music levels, and crowd noise. One study I have been involved in has evaluated the health effects of exposure to wind-turbine noise [5].

In legal liability issues and in workers' compensation board decisions, noise dosimetry is an essential tool in evidence.

I offer my appreciation to the occupational health and safety pioneers for developing a practical approach to hearing-loss risk assessment. These schemes have likely prevented more noise-induced hearing loss than any other policy.

References

1. HARRISON RV. (2008). Noise-induced hearing loss in children: A 'less than silent' environmental danger, *Paediatrics & Child Health*, Volume 13, Issue 5, May/June 2008, 377–382. <https://doi.org/10.1093/pch/13.5.377>
2. HARRISON RV. (2012) The prevention of noise induced hearing loss in children. *International Journal Pediatrics*, 2012 Article ID 473541, pp13 doi: 10.1155/2012/473541 <https://doi.org/10.1155/2012/473541>
3. Occupational Safety and Health Administration (OSHA). Noise exposure standards <https://www.osha.gov/noise/standards>
4. Canadian Centre for Occupational Health and Safety (2025). Noise, occupational Exposure limits in Canada. https://www.ccohs.ca/oshanswers/phys_agents/noise/exposure_can.html
5. Understanding the Evidence: Wind Turbine Noise. Ottawa: The Expert Panel on Wind Turbine Noise and Human Health, Council of Canadian Academies. ISBN 978-1-926522-07-4