

The Official Publication of the Canadian Academy of Audiology

Why is Tinnitus Hard to Measure?

Published June 28th, 2019

Brandon T. Paul, PhD



How do you "measure" someone's tinnitus, and what does it mean to do so? Clearly, quantifying an internal perception such as a phantom sound is not a trivial task, especially when there is no clear cause. And it may be unsurprising to learn that there is no consensus on how tinnitus should

be measured.¹ Despite this challenge, tinnitus measurement is a critical part of both clinical and neuroscientific inquiry. It helps scientists understand possible etiologies, as well as brain correlates that may explain the neural origins of tinnitus. Further, knowing which aspects of tinnitus are most bothersome for a person could help to direct possible treatments or coping strategies.

Tinnitus is most commonly measured using psychoacoustic methods that estimate or modulate the "acoustic properties" of tinnitus, or by self-report scales and questionnaires. Psychoacoustic

methods² present sounds that are matched in pitch or loudness level to a person's tinnitus. Masking noises can also be used to determine the level at which tinnitus is no longer perceived, or to produce a temporary abatement of tinnitus (i.e., residual inhibition). These procedures are easily performed on a standard audiometer and sometimes use similar methods to those used in obtaining thresholds. Self-report measures, on the other hand, use rankings or numbering to estimate these acoustic properties or its impact on the individual. These range from simple 10-point visual analog scales that rank the severity, loudness, or annoyance associated with tinnitus, or questionnaires with subscales developed to capture secondary effects such as sleep problems, social support, depression, and other emotional effects. Common questionnaires include the Tinnitus Handicap

Inventory (THI)³ and the Tinnitus Functional Index (TFI).⁴ There are also ongoing attempts to use

brain imaging to detect the presence and degree of tinnitus,^{5–7} but more work is needed before these tests are considered viable objective measures.

Despite the development of these tools, numerous factors – far too many to list here – lead to basic errors in tinnitus measurement. For psychoacoustic tasks, measurement reliability is reduced by factors including the presence of hearing loss that can make pitches hard to discriminate, cases of "complex" tinnitus that contain many concurrent pitches or noises, and the fact that tinnitus

loudness and pitch can change over time.^{2,8} A common problem with self-report measures is the ambiguity in what an individual should be rating. For example, it is common to ask a person to rate

if tinnitus is of "high" or "low severity" from 1 to 10.² Is it severe in loudness, annoyance, or other effects on everyday life? This may depend on what parts of the tinnitus experience, at the time of measurement, are most salient to the affected individual.

Aside from these sources of error, there are a handful of broader, more complex factors that may better explain systematic difficulties in tinnitus measurement. An obvious consideration is that tinnitus is associated with widespread changes in the brain, far beyond basic auditory perception. Neuroimaging studies routinely find that individuals with tinnitus have different levels of activity in brain areas involved in general neural function, such as attention, memory, and emotion, when

compared to non-tinnitus individuals.⁹⁻¹¹ Although these findings are unsurprising given the effects of tinnitus more broadly on quality of life, one important observation from scientific research is that the psychological impacts of tinnitus are often not well predicted by the acoustic properties of

tinnitus.² One commonly reported finding is that the loudness of someone's tinnitus estimated by psychoacoustic matching does not always correlate well to subjective ratings of annoyance,

distress, or even the self-reported rating of loudness itself.^{12,13} Although this is perplexing, we must consider that just because tinnitus is not comparatively loud does not mean that the perception is not pervasive, hard to ignore, and resistant to masking. This notion has important implications, and can confound tinnitus measurements. For instance, a person may expect the loudness of their tinnitus is about a "7" on a scale of 0 (no sound) to 10 (loudest imaginable), but when tinnitus loudness is matched to an external tone, it is below 10 dB sensation level (very low). The rated loudness level can be confounded by how strongly an individual is affected by their tinnitus. Thus, it is critical to consider that the auditory experience of tinnitus can be separate from its impact on the individual, and this highlights the importance of using a combination of different measurements.

Another complexity affecting tinnitus measurement is the heterogeneity across the tinnitus

population.¹⁴ Tinnitus can arise from a variety of etiologies, is mildly bothersome for some but catastrophic to others, and is associated with widely differing effects on hearing, cognition, and daily life function. Teasing apart these effects is not straightforward. For instance, does a high distress or severity rating directly reflect the impact of tinnitus, or does it reflect common

comorbidities such reduced sound level tolerance, hearing loss, or dizziness?¹⁵ Unfortunately, charting all factors contributing to the heterogeneity of tinnitus is daunting and is likely not

feasible using a single method (note that some have tried¹⁶). However, what may make tinnitus "severe" may depend on a specific configuration of underlying effects, and likely differs across individuals.

Overall, tinnitus measurement is difficult owing to the complexity of its experience. Although psychoacoustic methods are useful when approximating the sound qualities of tinnitus, they do not reliably predict the impact of tinnitus on the individual. The impacts of tinnitus are furthermore

hard to disentangle from those related to other auditory, cognitive, and personality factors. Nonetheless, quantifying the tinnitus experience is crucial for developing evidence-based treatments and intervention strategies, and should be encouraged in audiological practice.

Although there is no widely effective treatment for tinnitus, measurement can be considered the first step. The act of quantifying tinnitus and simply providing a "result" on paper can help a tinnitus sufferer validate their condition, and in itself is therapeutic.

References

- 1. Henry JA. "Measurement" of tinnitus. Otol Neurotol 2016 Sep 1;37(8):e276-85.
- 2. Henry JA, Meikle MB. Psychoacoustic measures of tinnitus. J Am Acad Audiol 2000 Mar 1;11(3):138–55.
- 3. Newman CW, Jacobson GP, Spitzer JB. Development of the tinnitus handicap inventory. Arch Otolaryngol Head Neck Surg 1996 Feb 1;122(2):143–8.
- 4. Meikle MB, Henry JA, Griest SE, et al. The tinnitus functional index: development of a new clinical measure for chronic, intrusive tinnitus. Ear Hear 2012 Mar 1;33(2):153–76.
- Paul BT, Schoenwiesner M, Hébert S. Towards an objective test of chronic tinnitus: Properties of auditory cortical potentials evoked by silent gaps in tinnitus-like sounds. Hearing Res 2018 Sep 1;366:90–8.
- 6. Han JH, Won JY, Hong SK, et al. Objective measurement of subjective tinnitus using the acoustic change complex. PloS One 2017 Nov 27;12(11):e0188268.
- Jackson R, Vijendren A, Phillips J. Objective measures of tinnitus: a systematic review. Otol Neurotol 2019 Feb 1;40(2):154–63.
- 8. Meikle MB. The interaction of central and peripheral mechanisms in tinnitus. Mechan Tinnitus 1995:181–206.
- 9. Roberts LE, Husain FT, Eggermont JJ. Role of attention in the generation and modulation of tinnitus. Neurosci Biobehavior Rev 2013 Sep 1;37(8):1754–73.
- 10. De Ridder D, Elgoyhen AB, Romo R, Langguth B. Phantom percepts: tinnitus and pain as persisting aversive memory networks. Proc Nat Acad Sci 2011 May 17;108(20):8075–80.
- 11. Husain FT. Neural networks of tinnitus in humans: elucidating severity and habituation. Hear Res 2016 Apr 1;334:37–48.
- 12. Fowler EP. The "illusion of loudness" of tinnitus—its etiology and treatment. Laryngoscope 1942 Apr;52(4):275–85.
- Hall DA, Mehta RL, Fackrell K. How to choose between measures of tinnitus loudness for clinical research? A report on the reliability and validity of an investigator-administered test and a patient-reported measure using baseline data collected in a phase IIa drug trial. Am J Audiol 2017 Sep 18;26(3):338–46.
- Cederroth CR, Gallus S, Hall DA, et al. Towards an understanding of tinnitus heterogeneity. Frontiers Aging Neurosci 2019;11:53.
- 15. Ratnayake SA, Jayarajan V, Bartlett J. Could an underlying hearing loss be a significant factor in the handicap caused by tinnitus? Noise Health 2009 Jul 1;11(44):156.
- 16. Schlee W, Hall DA, Edvall NK, et al. Visualization of global disease burden for the optimization of patient management and treatment. Frontiers Med 2017 Jun 19;4:86.